

Tumour budding (Non-core)

Tumour budding is defined as single cells or clusters of up to four tumour cells at the invasive front of carcinomas. It is considered to be the morphological manifestation of epithelial mesenchymal transition.¹ Tumour budding is different from tumour grade (based on gland formation) and poorly differentiated clusters (≥ 5 cells).

There is increasing evidence that tumour budding is an independent adverse prognostic factor in colorectal carcinoma. Several studies have shown that stage pT1 colorectal carcinomas with tumour budding score Bd2 and Bd3 (≥ 5 buds) are associated with an increased risk of lymph node metastasis.²⁻⁶ For stage II colorectal carcinomas, tumour budding score Bd3 is associated with increased risk of recurrence and mortality.⁷⁻⁹

Tumour budding is reported as the number of buds and scored using a three-tier system. According to the recommendations from a consensus conference on tumour budding,¹⁰ the number of tumour buds is the highest count after scanning ten separate fields (at 20x objective lens) along the invasive front of the tumour or the entire lesion for malignant polyps ('hotspot' approach). The number of tumour buds is counted on haematoxylin and eosin (H&E). If the invasive front of the tumour is obscured by inflammatory cells, immunohistochemistry using pan-cytokeratin can be used to help identifying the buds, but the final count is performed on H&E. Depending on the eyepiece field diameter of the microscope, the number of buds may need to be normalised to represent the number for a field of 0.785 mm² (objective lens 20x with eyepiece diameter of 20 mm).

Tumour budding should only be reported in non-mucinous and non-signet-ring cell adenocarcinoma areas. Furthermore, for colonic or rectal adenocarcinomas resected after neoadjuvant therapy, tumour budding should not be reported.

References

- 1 Zlobec I and Lugli A (2010). Epithelial mesenchymal transition and tumor budding in aggressive colorectal cancer: tumor budding as oncotarget. *Oncotarget* 1(7):651-661.
- 2 Pai RK, Cheng YW, Jakubowski MA, Shadrach BL, Plesec TP and Pai RK (2017). Colorectal carcinomas with submucosal invasion (pT1): analysis of histopathological and molecular factors predicting lymph node metastasis. *Mod Pathol* 30(1):113-122.
- 3 Beaton C, Twine CP, Williams GL and Radcliffe AG (2013). Systematic review and meta-analysis of histopathological factors influencing the risk of lymph node metastasis in early colorectal cancer. *Colorectal Dis* 15(7):788-797.
- 4 Ueno H, Hase K, Hashiguchi Y, Shimazaki H, Yoshii S, Kudo SE, Tanaka M, Akagi Y, Suto T, Nagata S, Matsuda K, Komori K, Yoshimatsu K, Tomita Y, Yokoyama S, Shinto E, Nakamura T and Sugihara K (2014). Novel risk factors for lymph node metastasis in early invasive colorectal cancer: a multi-institution pathology review. *J Gastroenterol* 49(9):1314-1323.
- 5 Suh JH, Han KS, Kim BC, Hong CW, Sohn DK, Chang HJ, Kim MJ, Park SC, Park JW, Choi HS and Oh JH (2012). Predictors for lymph node metastasis in T1 colorectal cancer. *Endoscopy* 44(6):590-595.

- 6 Tateishi Y, Nakanishi Y, Taniguchi H, Shimoda T and Umemura S (2010). Pathological prognostic factors predicting lymph node metastasis in submucosal invasive (T1) colorectal carcinoma. *Mod Pathol* 23(8):1068-1072.
- 7 van Wyk HC, Park J, Roxburgh C, Horgan P, Foulis A and McMillan DC (2015). The role of tumour budding in predicting survival in patients with primary operable colorectal cancer: a systematic review. *Cancer Treat Rev* 41(2):151-159.
- 8 Petrelli F, Pezzica E, Cabiddu M, Coinu A, Borgonovo K, Ghilardi M, Lonati V, Corti D and Barni S (2015). Tumour budding and survival in stage II colorectal cancer: a systematic review and pooled analysis. *J Gastrointest Cancer* 46(3):212-218.
- 9 Rogers AC, Winter DC, Heeney A, Gibbons D, Lugli A, Puppa G and Sheahan K (2016). Systematic review and meta-analysis of the impact of tumour budding in colorectal cancer. *Br J Cancer* 115(7):831-840.
- 10 Lugli A, Kirsch R, Ajioka Y, Bosman F, Cathomas G, Dawson H, El Zimaity H, Flejou JF, Hansen TP, Hartmann A, Kakar S, Langner C, Nagtegaal I, Puppa G, Riddell R, Ristimaki A, Sheahan K, Smyrk T, Sugihara K, Terris B, Ueno H, Vieth M, Zlobec I and Quirke P (2017). Recommendations for reporting tumor budding in colorectal cancer based on the International Tumor Budding Consensus Conference (ITBCC) 2016. *Mod Pathol* 30(9):1299-1311.