

# Lymphatic and venous invasion (Core)

For colorectal cancer, it is important to report the presence or absence of lymphovascular invasion and to classify this further according to the type of vessels involved and, for veins, their intramural or extramural location, as these features may have different clinical and prognostic implications.

Small vessel invasion should be reported separately from venous (large vessel) invasion. Small vessel invasion is defined as tumour involvement of thin-walled structures lined by endothelium, without an identifiable smooth muscle layer or elastic lamina. Small vessels may represent lymphatics, capillaries or post-capillary venules. Lymphatics and venules may be distinguished by D2-40 immunohistochemistry, which only stains lymphatic endothelial cells, not venular, but this is not routinely recommended in reporting polypectomy specimens. All forms of small vessel invasion are considered under the 'L' classification under Union for International Cancer Control (UICC)/American Joint Committee on Cancer (AJCC) TNM 8<sup>th</sup> editions.<sup>1,2</sup> Small vessel invasion is associated with lymph node metastatic disease and has been shown to be an independent indicator of adverse outcome in several studies.<sup>3-5</sup> A higher prognostic significance of extramural small vessel invasion has been suggested, but the importance of anatomic location in small vessel invasion is not well established.<sup>4</sup> In malignant polyps, lymphatic invasion is associated with an increased risk of lymph node metastasis.<sup>6,7</sup>

Venous invasion is defined as tumour present within an endothelium-lined space that is either surrounded by a rim of muscle or contains red blood cells.<sup>8</sup> It should be suspected in the presence of a rounded or elongated deposit of tumour beside an artery. Interpretation of such features is subjective and can be improved by the application of immunohistochemical and histochemical stains, in particular elastic staining to identify venous elastic lamina.<sup>9,10</sup> A circumscribed tumour nodule surrounded by a smooth muscle wall or an identifiable elastic lamina, evident on haematoxylin & eosin (H&E) or elastic stains, is considered sufficient to classify as venous invasion. Examination of multiple levels in blocks showing features suspicious of venous invasion can also be helpful in borderline cases.

Extramural (beyond muscularis propria) venous invasion has been demonstrated by multivariate analysis in multiple studies to be a stage independent adverse prognostic factor for colonic and rectal cancer.<sup>4,11</sup> However, it would only be identified in transanal minimally invasive surgery (TAMIS) or endoscopic full thickness resection (EFTR) specimens. There is also evidence from several studies that intramural (intramuscular or submucosal) venous spread is also of prognostic importance but the evidence is much weaker than for extramural venous invasion.<sup>12</sup>

## References

- 1 Brierley JD, Gospodarowicz MK and Wittekind C (eds) (2016). *UICC TNM Classification of Malignant Tumours, 8th Edition*, Wiley-Blackwell.
- 2 Amin MB, Edge SB, Greene FL, Byrd DR, Brookland RK, Washington MK, Gershenwald JE, Compton CC, Hess KR, Sullivan DC, Jessup JM, Brierley JD, Gaspar LE, Schilsky RL, Balch CM, Winchester DP, Asare EA, Madera M, Gress DM and Meyer LR (eds) (2017). *AJCC Cancer Staging Manual. 8th ed.*, Springer, New York.
- 3 Santos C, Lopez-Doriga A, Navarro M, Mateo J, Biondo S, Martinez Villacampa M, Soler G, Sanjuan X, Paules MJ, Laquente B, Guino E, Kreisler E, Frago R, Germa JR, Moreno V and Salazar R (2013). Clinicopathological risk factors of Stage II colon cancer: results of a prospective study. *Colorectal Dis* 15(4):414-422.

- 4 Betge J, Pollheimer MJ, Lindtner RA, Kornprat P, Schlemmer A, Rehak P, Vieth M, Hoeffler G and Langner C (2012). Intramural and extramural vascular invasion in colorectal cancer: prognostic significance and quality of pathology reporting. *Cancer* 118(3):628-638.
- 5 Lim SB, Yu CS, Jang SJ, Kim TW, Kim JH and Kim JC (2010). Prognostic significance of lymphovascular invasion in sporadic colorectal cancer. *Dis Colon Rectum* 53(4):377-384.
- 6 Beaton C, Twine CP, Williams GL and Radcliffe AG (2013). Systematic review and meta-analysis of histopathological factors influencing the risk of lymph node metastasis in early colorectal cancer. *Colorectal Dis* 15(7):788-797.
- 7 Hassan C, Zullo A, Risio M, Rossini FP and Morini S (2005). Histologic risk factors and clinical outcome in colorectal malignant polyp: a pooled-data analysis. *Dis Colon Rectum* 48(8):1588-1596.
- 8 Talbot IC, Ritchie S, Leighton M, Hughes AO, Bussey HJ and Morson BC (1981). Invasion of veins by carcinoma of rectum: method of detection, histological features and significance. *Histopathology* 5(2):141-163.
- 9 Howlett CJ, Tweedie EJ and Driman DK (2009). Use of an elastic stain to show venous invasion in colorectal carcinoma: a simple technique for detection of an important prognostic factor. *J Clin Pathol* 62(11):1021-1025.
- 10 Kirsch R, Messenger DE, Riddell RH, Pollett A, Cook M, Al-Haddad S, Streutker CJ, Divaris DX, Pandit R, Newell KJ, Liu J, Price RG, Smith S, Parfitt JR and Driman DK (2013). Venous invasion in colorectal cancer: impact of an elastin stain on detection and interobserver agreement among gastrointestinal and nongastrointestinal pathologists. *Am J Surg Pathol* 37(2):200-210.
- 11 Freedman LS, Macaskill P and Smith AN (1984). Multivariate analysis of prognostic factors for operable rectal cancer. *Lancet* 2(8405):733-736.
- 12 Knijn N, van Exsel UEM, de Noo ME and Nagtegaal ID (2018). The value of intramural vascular invasion in colorectal cancer - a systematic review and meta-analysis. *Histopathology* 72(5):721-728.