


# Neoplasia of the Testis - Retroperitoneal Lymphadenectomy

## Histopathology Reporting Guide

Family/Last name

Given name(s)  Date of birth

Patient identifiers  Date of request  Accession/Laboratory number

Elements in **black text** are REQUIRED. Elements in **grey text** are RECOMMENDED. SCOPE OF REPORTING GUIDE 

### CLINICAL INFORMATION

Not provided

Previous history of testicular cancer, *specify*

Previous therapy, *specify*

Other, *specify*

### PRE-PROCEDURE SERUM TUMOUR MARKERS

(select all that apply)

Not provided

Provided

Serum tumour markers within normal limits OR Specify serum tumour markers used, level and date markers were drawn

Date   AFP

LDH   b-HcG

### SPECIMEN(S) SUBMITTED (select all that apply)

Not specified

Retroperitoneal lymphadenectomy, *specify nodal site*

No disease

Necrosis

Viable tumour

No disease

Necrosis

Viable tumour

No disease

Necrosis

Viable tumour

Brain  Liver

Lung  Other, *specify*

### SIZE OF LARGEST NODAL METASTASIS (select all that apply)

Cannot be assessed

Maximum dimension

mm

Additional dimensions

mm X  mm

### BLOCK IDENTIFICATION KEY

(List overleaf or separately with an indication of the nature and origin of all tissue blocks)

### HISTOLOGICAL TUMOUR TYPE

(Value list from the World Health Organisation Classification of tumours. Pathology and genetics of urinary system and male genital organs (2016))

Viable tumour

Present  Absent

%

Germ cell tumour, *specify type and percentage*

→  %

→  %

→  %

→  %

Other, *specify*

### MARGIN STATUS

Cannot be assessed

Not involved

Closest margin

Distance of tumour from closest margin  mm

Involved, *specify*

## EXTRANODAL EXTENSION

- Not identified     Present     Indeterminate

## PATHOLOGIC STAGING (TNM 8th edition)\*\*

### Regional lymph nodes (pN)

- NX    Regional lymph nodes cannot be assessed
- N0    No regional lymph node metastasis
- N1    Metastasis with a lymph node mass 2 cm or smaller in greatest dimension and less than or equal to five nodes positive, none larger than 2 cm in greatest dimension
- N2    Metastasis with a lymph node mass larger than 2 cm but not larger than 5 cm in greatest dimension; or more than five nodes positive, none larger than 5 cm; or evidence of extranodal extension of tumour
- N3    Metastasis with a lymph node mass larger than 5 cm in greatest dimension

### Distant metastasis (pM) (if resected)

- No distant metastases
- M1    Distant metastasis
- M1a    Non-retroperitoneal nodal or pulmonary metastases
- M1b    Non-pulmonary visceral metastases

##    Used with the permission of the American College of Surgeons, Chicago, Illinois. The original source for this information is the AJCC Cancer Staging Manual, Eighth Edition (2016) published by Springer Science+Business Media.

^    Please note that implementation of AJCC TNM 8th edition has been deferred until January 2018 in some jurisdictions. UICC 7th edition or AJCC 7th edition may be useful in the interim.