

Prostate Cancer Histopathology Reporting Guide Radical Prostatectomy Specimen



Radical Prosta	atectomy Specimen
Family/Last name	Date of birth DD - MM - YYYY
Given name(s)	
Patient identifiers	Date of request Accession/Laboratory number
	DD - MM - YYYY
Elements in black text are CORE. Elements in grey text are N indicates multi-select values indicates single select values	SCOPE OF THIS DATASET
CLINICAL INFORMATION	LYMPH NODE DISSECTION SPECIMEN(S)
Information not provided	Not submitted
Information provided (select all that apply)	Present (partially or completely resected)
Previous history of prostate cancer (including the Gleason score or WHO/ISUP Grade/Grade Group of previous specimens if known), specify	Site(s), specify
	Laterality
	Left Right Bilateral Other
Previous biopsy, specify date and where performed	
Trevious biopsy, speeny date and where performed	BLOCK IDENTIFICATION KEY (List overleaf or separately with an indication of the nature and origin of all tissue blocks)
Previous therapy, specify	
	HISTOLOGICAL TUMOUR TYPE (select all that apply)
	Adenocarcinoma (Acinar, usual type) Other, <i>specify</i>
Other clinical information, specify	
PRE-BIOPSY SERUM PSA	HISTOLOGICAL TUMOUR GRADE
	Gleason score
ng/mL	Indicate how Gleason score is being reported Largest tumour nodule
	Highest score tumour nodule
SPECIMEN WEIGHT (Weight of the prostate gland without the seminal vesicles)	Highest pT category tumour nodule
(weight of the prostate gland without the seminal vesicles)	Global score (summation of Gleason patterns in all floudies)
g	Primary pattern
	Secondary pattern
SPECIMEN DIMENSIONS (Of the prostate gland)	
(or the prostate giant)	Indeterminate, specify reason
length mm x width mm x depth mm	
SEMINAL VESICLES	
Absent	Minor tertiary pattern (if present and higher than primary and secondary grade)
Present (partially or completely resected)	4 5 Not applicable

WHO/ISUP Grade (Grade Group) WHO/ISUP Grade (Grade Group) 1 (Gleason score ≤6) WHO/ISUP Grade (Grade Group) 2 (Gleason score 3+4=7) WHO/ISUP Grade (Grade Group) 3 (Gleason score 4+3=7) WHO/ISUP Grade (Grade Group) 4 (Gleason score 8) WHO/ISUP Grade (Grade Group) 5 (Gleason score 9-10) Indeterminate, specify reason	MICROSCOPIC URINARY BLADDER NECK INVASION Not applicable Not identified Present Refers to cases where bladder neck is not included in the specimen. MARGIN STATUS Cannot be assessed Not involved Involved, specify margin(s) and their location, if possible
TUMOUR GROWTH PATTERNS	
Intraductal carcinoma of the prostate (IDC-P) AND/OR Invasive cribriform carcinoma (ICC) Indeterminate Not identified Present If present, specify the tumour growth pattern (if apparent	Type of margin positivity (select all that apply) Indeterminate Extraprostatic (EPE) Intraprostatic (capsular incision)
on H&E staining ^a)	Length of margin involved by carcinoma ^c
IDC-P Not identified Present	mm
IDC-P incorporated into Gleason score	Gleason pattern of tumour present at positive margin
☐ IDC-P not incorporated into Gleason score	○ Gleason pattern 3
Invasive cribriform carcinoma	☐ Gleason pattern 4 or/and 5
Not identified	^c If more than 1 positive margin the extent should reflect the cumulative
Present	length.
^a Use of immunohistochemistry is optional.	^d If more than 1 pattern at margin select the highest.
TUMOUR QUANTIFICATION (Amount of tumour identified)	SEMINAL VESICLE INVASION
Percentage of prostate involved by tumour	 Not applicable^e Not identified Present Refers to rare cases where seminal vesicles are not included in the specimen.
	LYMPHOVASCIII AD INVASTONI E
OR Diameter of largest nodule	LYMPHOVASCULAR INVASION☐ Indeterminate☐ Not identified☐ Present
	LYMPH NODE STATUS
EXTRAPROSTATIC EXTENSION	No nodes submitted or found
○ Indeterminate○ Not identified	Number of lymph nodes examined
Present	○ Not involved
Specify location(s)	Involved
	Number of involved lymph nodes
	Number cannot be determined
Extent Focal Non-focal (established)	Maximum dimension of largest mm

PATHOLOG:	ICAL STAGING (UICC TNM 8 th edition) ^f	
	riptors (only if applicable) (select all that apply)	
□ m -	multiple primary tumours	
□ ''' □ r -	recurrent	
□ ·	post neoadjuvant therapy	
_ ,		
	ımour (pT)	
\bigcirc TX ⁹	Primary tumour cannot be assessed	
○ T0	No evidence of primary tumour	
	Tumour confined within prostate	
○ T3	Tumour extends through the prostatic capsule ^{h,i}	
	Extracapsular extension ^h (unilateral or bilateral) including microscopic bladder neck involvement	
	Tumour invades seminal vesicle(s)	
○ T4	Tumour is fixed or invades adjacent structures other than seminal vesicles: external sphincter, rectum, levator muscles and/or pelvic wall	
Regional lymph nodes (pN)		
\bigcirc NX ^g	Regional nodes cannot be assessed	
○ N0	No regional lymph node metastasis	
○ N1	Regional lymph node metastasis	
Distant me	etastasis (pM) ^j	
	pplicable ^k	
○ M1	Distant metastasis	
\circ	Non-regional lymph node(s)	
	Bone(s)	
	c Other site(s)	
0 1120		
Malignant Tui Gospodarowid	with permission. Source: UICC TNM Classification of mours, 8 th Edition, eds by James D. Brierley, Mary K. cz, Christian Wittekind. 2016, Publisher Wiley. g any errata published up until 12 th July 2024).	
TX and NX sh	nould be used only if absolutely necessary.	
The consensus of the dataset authors is that the term extraprostatic extension is preferred.		
Invasion into the prostatic apex or into (but not beyond) the prostatic capsule is not classified as T3, but as T2.		
	more than 1 site of metastasis is present, the most egory is used. pM1c is the most advanced category.	
< No clinical an	d radiological correlation available.	
ciiincai aii	g.ca. co classon aranables	